

学校编码: 10384

分类号_____密级_____

学号: 20520130153870

UDC_____

厦 门 大 学

博 士 学 位 论 文

负载钌、钯催化剂上乙酰丙酸及
对乙酰氨基苯酚选择加氢的研究

Studies on Selective Hydrogenation of Levulinic Acid and
Paracetamol over Supported Ruthenium and Palladium
Catalysts

宋 文 静

指导教师姓名: 伊晓东 副教授

翁维正 教 授

专业名称: 物 理 化 学

论文提交日期: 2016 年 08 月

论文答辩日期: 2016 年 09 月

学位授予日期: 2016 年 月

答辩委员会主席: _____

评 阅 人: _____

2016 年 09 月

The Doctor Dissertation of Science in Xiamen University

**Studies on Selective Hydrogenation of Levulinic Acid and
Paracetamol over Supported Ruthenium and Palladium
Catalysts**

Wenjing Song

Supervised by

Professors: Xiaodong Yi

Weizheng Weng

Department of Chemistry

Xiamen University

September, 2016

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文(包括纸质版和电子版)，允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

()1.经厦门大学保密委员会审查核定的保密学位论文，于
年 月 日解密，解密后适用上述授权。

()2.不保密，适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。)

声明人(签名)：

年 月 日

目 录	
摘 要.....	I
Abstract.....	III
第一章 绪论	1
1.1 乙酰丙酸加氢制戊内酯的研究	1
1.1.1 研究意义.....	3
1.1.2 均相催化剂.....	5
1.1.3 多相催化剂.....	6
1.1.4 乙酰丙酸加氢机理研究.....	14
1.2 对乙酰氨基苯酚选择加氢制酮的研究	15
1.2.1 苯环加氢反应体系.....	17
1.2.2 苯酚加氢反应体系.....	19
1.2.3 取代苯酚加氢反应体系.....	23
1.2.4 对乙酰氨基苯酚加氢反应体系.....	25
1.3 论文的构思与目的	25
1.3.1 乙酰丙酸加氢制戊内酯.....	25
1.3.2 对乙酰氨基苯酚加氢制对乙酰氨基环己酮.....	26
1.4 论文的组成与概要	27
参考文献	28
第二章 实验部分	44
2.1 主要气体和试剂	44
2.2 催化剂常规表征	45
2.2.1 X-射线粉末衍射 (XRD)	45
2.2.2 N ₂ 吸脱附	45
2.2.3 程序升温还原实验 (H ₂ -TPR)	46
2.2.4 热重分析 (TG)	46
2.2.5 X 射线光电子能谱 (XPS)	46

2.2.6 激光 Raman 光谱	47
2.2.7 透射电子显微镜 (TEM)	47
2.2.8 CO ₂ 程序升温脱附 (CO ₂ -TPD)	47
2.2.9 H ₂ 程序升温脱附 (H ₂ -TPD)	48
2.2.10 CO 化学吸附测试	48
2.2.11 吡啶-红外分析测试 (Py-IR)	48
参考文献	49
第三章 不同碳材料负载钌催化剂上乙酰丙酸加氢反应的研究	50
3.1 引言	50
3.2 反应原理	51
3.3 实验部分	52
3.3.1 催化剂制备	52
3.3.2 催化剂性能评价	54
3.4 结果与讨论	55
3.4.1 不同载体负载 Ru 基催化剂的液相乙酰丙酸加氢性能	55
3.4.2 不同反应条件下液相乙酰丙酸加氢性能	59
3.4.3 Ru/HSAG500 催化剂的乙酰丙酸加氢稳定性考察	61
3.4.4 催化剂的表征及构效关联	62
3.5 本章小结	76
参考文献	77
第四章 La 助剂修饰 Ru/CF 催化剂上乙酰丙酸加氢的研究	85
4.1 引言	85
4.2 实验部分	87
4.2.1 催化剂制备	87
4.2.2 催化剂性能评价	87
4.3 结果与讨论	87
4.3.1 助剂种类对 Ru/CF 催化性能的影响	87
4.3.2 助剂 La 添加量对 Ru/C 催化性能的影响	89
4.3.3 Ru _x La/CF 催化剂表征	91

4.3.4 Ru _x La/CF 催化剂反应动力学实验及模型拟合计算.....	104
4.3.5 Ru _x La/CF 催化剂的结构与性能关联.....	111
4.4 本章小结	114
参考文献	115
第五章 硅烷化改性 SiO₂ 负载 Pd 催化剂的选择加氢反应性能.....	121
5.1 引言	121
5.2 反应机理	122
5.3 实验部分	123
5.3.1 催化剂制备.....	123
5.3.2 催化剂性能评价.....	124
5.4 结果与讨论	125
5.4.1 二氧化硅负载型钯基催化剂的对乙酰氨基苯酚加氢性能.....	125
5.4.2 硅烷化处理催化剂对乙酰氨基苯酚加氢性能的影响.....	131
5.4.3 负载型钯基催化剂的苯酚加氢性能.....	135
5.4.4 硅烷化后催化剂的表征及构效关联.....	136
5.5 本章小结	145
参考文献	145
第六章 总结	151
6.1 不同碳材料负载钨催化剂上乙酰丙酸加氢反应的研究	151
6.2 La 助剂修饰 Ru/CF 催化剂上乙酰丙酸加氢的研究	152
6.3 硅烷化改性 SiO ₂ 负载 Pd 催化剂的选择加氢反应性能	153
在研期间发表论文及专利	154
致 谢.....	156

Contents

Abstract in Chinese.....	I
Abstract in English	III
Chapter 1 Introduction.....	1
1.1 Hydrogenation of levulinic acid to γ-valerolactone.....	1
1.1.1 The significance of the research	3
1.1.2 The homogeneous catalyst.....	5
1.1.3 The hetrogeneous catalyst.....	6
1.1.4 Mechanistic aspects of hydrogenation of levulinic acid	14
1.2 Hydrogenation of paracetamol to acetamidocyclohexanone	15
1.2.1 Selective hydrogenation of benzene	17
1.2.2 Selective hydrogenation of phenol.....	19
1.2.3 Selective hydrogenation of substituted phenols.....	23
1.2.4 Selective hydrogenation of paracetamol	25
1.3 Objectives of this thesis	25
1.3.1 Hydrogenation of levulinic acid to γ -valerolactone.....	25
1.3.2 Hydrogenation of paracetamol to acetamidocyclohexanone	26
1.4 Outline of this thesis.....	27
References.....	28
Chapter 2 Experimental	44
2.1 Materials and reagents	44
2.2 Charaterizations of catalysts.....	45
2.2.1 X-ray diffraction (XRD)	45
2.2.2 N ₂ physical adsorption (BET).....	45
2.2.3 H ₂ temperature programmed reduction (H ₂ -TPR)	46
2.2.4 Thermogravimetry (TG)	46
2.2.5 X-ray phtoelectron spectroscopy (XPS)	46

2.2.6 Laser Raman spectroscopy	47
2.2.7 Transmission electron microscopy (TEM).....	47
2.2.8 CO ₂ temperature programmed decomposition (CO ₂ -TPD)	47
2.2.9 H ₂ temperature programmed decomposition (H ₂ -TPD)	48
2.2.10 CO chemisorption	48
2.2.11 Py-IR characterization.....	48
Reference	49
 Chapter 3 Different carbon supported Ru catalysts for levulinic acid	
hydrogenation.....	50
3.1 Introduction.....	50
3.2 Reaction mechanism	51
3.3 Experimental	52
3.3.1 Preparation of catalysts	52
3.3.2 Evaluation of catalytic properties	54
3.4 Results and discussions.....	55
3.4.1 Catalytic performances of Ru catalysts loaded on different supports for the hydrogenation of levulinic acid	55
3.4.2 Effect of reaction conditions on the LA hydrogenation	59
3.4.3 Stability of the Ru/HSAG500 catalysts	61
3.4.4 Chacterizations and structure-activity relationships	62
3.5 Conclusions.....	76
References	77
 Chapter 4 Effect of lanthanum promoter on the catalytic	
performance of levulinic acid hydrogenation over Ru/Carbon fiber	
catalyst	85
4.1 Introduction.....	85
4.2 Experimental	87
4.2.1 Preparation of catalysts	87

4.2.2 Evaluation of catalytic properties	87
4.3 Results and discussions.....	87
4.3.1 Effect of promoter on the catalytic performance	87
4.3.2 Effect of promoter loadings on catalytic performance	89
4.3.3 Characterizations of the RuxLa/CF catalysts.....	91
4.3.4 Kinetic test and fitting calculation	104
4.3.5 Correlation between the structure and the catalytic behaviours of RuxLa/CF.....	111
4.4 Conclusions.....	114
References.....	115
 Chapter 5 Selective hydrogenation of paracetamol to acetamidocyclohexanone with silylated SiO₂ supported Pd-based catalysts.....	 121
5.1 Introduction.....	121
5.2 Reaction mechanism	122
5.3 Experimental	123
5.3.1 Preparation of catalysts	123
5.3.2 Evaluation of catalytic properties	124
5.4 Results and discussions.....	125
5.4.1 Catalytic performances of Pd catalysts loaded on SiO ₂ for the hydrogenation of paracetamol.....	125
5.4.2 Catalytic performances of Pd catalysts loaded on silylated SiO ₂ for the hydrogenation of paracetamol.....	131
5.4.3 Catalytic behaviours of Pd catalysts for phenol hydrogenation.....	135
5.4.4 Correlation between the structure of silylated SiO ₂ supported Pd-based catalysts and the catalytic behaviours of paracetamol hydrogenation	136
5.5 Conclusions.....	145
References.....	145

Chapter 6 General conclusionss	151
6.1 Different carbon supported Ru catalysts for levulinic acid hydrogenation	151
6.2 Effect of lanthanum promoter on the catalytic performance of levulinic acid hydrogenation over Ru/Carbon fiber catalyst	152
6.3 Selective hydrogenation of paracetamol to acetamidocyclohexanone with silylated SiO₂ supported Pd-based catalysts.....	153
List of publications and patents.....	154
Acknowledgements	156

摘 要

本论文以重要的生物质平台化合物乙酰丙酸(LA)加氢制备 γ -戊内酯(GVL)以及对乙酰氨基苯酚选择加氢制备酮为目标,针对选择性加氢的控制,系统研究了碳材料负载钌催化剂上乙酰丙酸加氢性能及硅烷化改性 SiO_2 负载钯基催化剂上对乙酰氨基苯酚的加氢性能。通过对催化剂的结构和性质进行较为系统的表征并与其催化活性进行关联,取得了以下主要结果。

论文针对 LA 选择加氢制备 GVL 的研究,比较了不同载体,包括不同碳材料及无机氧化物负载钌催化剂的反应性能。实验结果显示,反应产物 GVL 选择性均较高,达到 99 % 以上,而载体对钌基催化剂的 LA 加氢活性具有显著影响。不同碳材料物化性质如比表面积、孔结构、电子性质及表面化学性质等都不同,对应催化剂的催化反应活性也不尽相同。因此,我们研究不同碳材料(石墨、氧化石墨、还原性氧化石墨烯、碳纳米管、碳纤维、活性炭)对 Ru 基催化剂 LA 加氢性能的影响。TEM、CO 化学吸附热等测试表明,当 Ru 金属颗粒尺寸为 1.2-2.9 nm 之间时,乙酰丙酸加氢反应是结构敏感反应, Ru 粒子尺寸越小, LA 加氢的 TOF 越高,且表面低配位的 Ru 有利于加氢反应的进行。氮气吸脱附实验及热重结果表明,载体的表面含氧官能团数量不同时,由于官能团与前驱盐之间的相互作用不同,影响金属颗粒大小,从而改变了 LA 加氢转化频率(TOF)。

另一方面,考虑到 La 在加氢反应中起到促进作用,而对 La 参与反应的机理研究仍然较少,我们将催化剂的 Ru 金属负载量降低为 1 %,提高 LA/Ru 底物比,在 100 °C、5 MPa 氢压反应条件下,考查了不同 La 助剂添加量对碳纤维负载 Ru 的 LA 加氢性能的影响。结果表明,La 含量为 0.5 % 的 $\text{Ru}_{0.5}\text{La}/\text{CF}$ 催化剂表现出最高的 GVL 收率,是未添加 La 助剂的 Ru/CF 催化剂的 2.6 倍。当 La 的添加过量时, LA 加氢活性有所下降。利用 CO 化学吸附、 CO_2 -TPD、 H_2 -TPR、XPS、TEM 等手段研究催化剂的结构和性质,表征结果显示,在 Ru 粒径大小基本一致(2 nm 左右)的情况下,La 物种负载在 Ru/C 表面后,降低了 Ru 金属的 CO 化学吸附量。La 物种与 Ru 纳米粒子之间的相互作用促使金属 Ru 电子富集,从而有利于电子反馈到 LA 的羰基轨道上,进而促进其加氢反应的进行。同时,富集在金属 Ru 周围的碱性 La 物种易吸附反应物乙酰丙酸,从而提升 Ru 活性位

对羰基的亲核进攻能力。催化剂表观加氢活性取决于 Ru 活性位点数目及由 Ru-La 界面相互作用等因素决定的 TOF 值。

由于在香料、医药等领域的重要应用，苯酚类化合物选择性加氢制备相应的酮、醇化合物一直以来都受到广泛关注。对乙酰氨基环己酮是合成麻醉药吗啡及噻唑类化合物，如治疗帕金森氏病的药物普拉克索等药物的重要中间体。论文针对乙酰氨基苯酚选择性加氢制备对乙酰氨基环己酮的反应，通过调变 Pd/SiO₂ 催化剂表面化学性质来改变其催化选择加氢性能。

以浸渍氨丙基三乙氧基硅烷 (APTES) 的方法来调变载体 SiO₂ 的表面结构，通过调控催化剂焙烧温度改变催化剂上硅烷化官能团的含量。TEM 结果表明，在 Pd 金属粒子大小为 4-7 nm 范围内，催化剂加氢活性与金属颗粒大小相关性低。硅烷化试剂 APTES 使载体表面存在一定量的官能团，而硅烷化官能团的含量影响对乙酰氨基苯酚加氢反应活性及酮的选择性。从 H₂-TPR 结果可知，由于催化剂焙烧温度不同，催化剂还原能力及 Pd-H 物种稳定性均有所不同。对于 500 °C 焙烧后的 Pd/Sily-500 催化剂，Pd-H 物种分解温度较低，H₂ 与 Pd 之间吸附较弱，易脱附的氢更有利于加氢反应的进行。当 H 原子在 Pd 表面难脱附时，不易参与加氢反应，导致催化活性较低。官能团的存在改变了 Pd 物种在载体表面的存在状态，硅烷化处理后 Pd 催化剂表面更易形成 PdH 物种，因此提高了 Pd/SiO₂ 催化剂的加氢活性。另一方面，对乙酰氨基苯酚通过苯环 π 电子结构吸附在催化剂 Pd 金属上进行反应，反应底物平面及非平面两种吸附方式影响了反应选择性的。硅烷化试剂的存在增强了反应的空间位阻效应，即表面官能团的存在阻碍了对乙酰氨基苯酚与催化剂之间通过 Pd-O 键的平面吸附，进而阻止了其进一步加氢反应，从而显著提高酮的选择性。对于未进行硅烷化处理的 Pd/SiO₂ 催化剂，反应物更倾向于以共平面吸附方式存在于催化剂表面，随后进行深度加氢生成对乙酰氨基环己醇。

关键词：乙酰丙酸；戊内酯；对乙酰氨基酚；对乙酰氨基环己酮；选择加氢

Abstract

This dissertation focused on the studies of the hydrogenation of levulinic acid (LA), a key platform compound from cellulose, into γ -valerolactone (GVL) and the selective hydrogenation of paracetamol into 4-acetamidocyclohexanone.

For the hydrogenation of LA, the catalytic performances of Ru nanoparticles loaded on the various supports were investigated in aqueous solutions, including the different oxide supports and carbon supports. It was found that the selectivity of GVL over Ru catalysts could reach 99 %, and the activity of the catalysts was affected significantly by their supports. Physicochemical properties of carbon materials, such as electrical conductivity, surface area and porosity, and chemical properties could be tuned for specific applications. Few reports deal with the influence of carbon characteristics such as nanostructure and surface chemical properties on the performance of the carbon supported Ru catalysts for the synthesis of γ -valerolactone from levulinic acid. Herein, the present work focuses on the Ru supported on very different carbon materials and their influences on the LA hydrogenation reaction. Ru-catalyzed hydrogenation of LA was a structure-sensitive reaction when the Ru particle size changed within 1.2-2.9 nm, small Ru nanoparticles favor the transformation of LA and the turnover frequency increased with decreasing the mean size of Ru nanoparticles. The test results of intrinsic activity and TOF in the LA hydrogenation reaction suggests that the low-coordination unsaturated Ru sites at the small metal nanoparticles are responsible for the improvement of catalytic activity. The characterizations by means of BET, TG were conducted and the results indicated that the amount of oxygen groups varied with support kinds, leading to a different Ru particle size. The differential heats of CO adsorption do not depend on the oxygen-contained groups, which were relevant to the particle sizes and electron density of ruthenium.

On the other hand, the incorporation of lanthanum on carbon nanofiber supported

Ru catalyst (a lower Ru loading) and its influence on the catalytic activity for the hydrogenation of LA to GVL was also studied in this work according the few study on the mechanism of the influence of La addition. Concerning the catalytic activity, at a temperature of 100 °C, hydrogen pressure of 5 MPa, the optimized Ru_{0.5}La/CF catalyst afforded 2.6 times GVL yield than the monometallic Ru catalyst. A maximum in the reaction rate ($53.8 \mu\text{mol}\cdot\text{gRu}^{-1} \text{ min}^{-1}$) is noted upon increasing La content (Ru_{0.5}La/CF), followed by a decline in catalytic activity at the lanthanum level in excess of the maximum. The satellite/main peak intensity ratio ($I_{\text{sat}}/I_{\text{mp}}$) from La 3d_{5/2} XPS spectra and TPR results indicated the presence of Ru interacting with La and the lanthanum was present in a different chemical environment. The shift of metallic Ru 3d peak positions to lower binding energy indicated the enhancement of electron density of Ru species by La addition to Ru/CF. Turnover frequencies for LA hydrogenation on RuLa/CF catalysts were found to increase with the addition of La mole fraction, suggesting the improvement of the surface Ru-La interfaces in the reactivity. Therefore, the catalytic performance was a consequence of both Ru metal which might be blocked by the dopant La and the Ru-La interfacial sites.

The hydrogenation of substituted phenols to the corresponding alkyl-substituted cyclohexanones and cyclohexanols has generated great interest in the industry and researchers, due to their application in the subsequent manufacture of fine chemicals for the fragrance and detergent industries. The 4-acetamidocyclohexanone was prepared as an intermediate in the synthesis of morphine skeleton. Nevertheless, reports on the produce of 4-acetamidocyclohexanone are still limited. Herein, in the present work, we attempt to tune the surface chemical composition of heterogeneous metal catalysts to optimize their hydrogenation performances.

For this, the silylated SiO₂ was prepared by impregnation of silica gel with the (3-aminopropyl) triethoxysilane (APTES) solution in toluene. The amount of sily groups on the SiO₂ support was tuned via the temperature of calcinations. TEM characterization indicated that the specific activity of catalyst did not seem to be greatly affected by the particle size considering the narrow size range of Pd particles (4-7 nm). From TPR results, the reducing ability of the catalysts and the stability of

the hydride phase could be tuned by the various calcination temperatures. The silylation process favors the formation of hydride phase on the Pd catalyst. Moreover, the palladium hydride species decompose at a lower temperature on the Pd/Sily-500, indicating a weaker absorption of H_2 on a Pd, which can facilitate the hydrogenation activity. Meanwhile, the APTES anchored on the support impedes the planar adsorption of paracetamol via O-Pd bonding on the catalyst surface due to steric hindrance, thus preventing them from further hydrogenation. The non-silylated SiO_2 supported Pd catalyst favors the co-planar absorption of paracetamol, obtaining the deeply hydrogenated product of 4-acetamidocyclohexanol.

Keywords: levulinic acid; γ -valerolactone; paracetamol; 4-acetamidocyclohexanone; selective hydrogenation

Degree papers are in the “[Xiamen University Electronic Theses and Dissertations Database](#)”.

Fulltexts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.